

SEG 100 Conference: Celebrating a Century of Discovery

ST.198

The Geometry and Kinematic History of the Howard's Pass XY Deposits

Darius Kamal, Ken Hickey

University of British Columbia, VANCOUVER, BC, Canada

The shale-hosted massive sulfide deposits of Howard's Pass are home to the world's most well-endowed undeveloped Zn-Pb resources. Sulfide deposition occurred in the Silurian, and the deposit was deformed during the Cretaceous Cordilleran orogeny. The Zn-Pb deposits have been incorporated into a series of upright WNW-striking folds on a 40-km trend along the border of the Yukon and Northwest Territories. Owing to the structural complexity of the deposits, multiple structural models for their evolution have been proposed. These range from Silurian soft-sediment deformation to the proposal of a regional shear zone hosting the deposits. In the shear zone model, bedding is completely transposed and previously interpreted stratigraphic boundaries are tectonic in origin. This current study aims to test this model and is focused on the XY group of deposits at the eastern end of the Howard's Pass district. Lithostratigraphic mapping and structural observations indicate one main phase of folding, F_1 , and the XY group of deposits is located on the southern limb of a macroscopic syncline. F_1 folds are upright and gently plunging to the WNW-NNW (Fig. 1). A regionally developed, steep NE-dipping cleavage, S_1 , is axial planar to the F_1 folds across Howard's Pass. S_1 typically manifests as a slaty cleavage comprising pervasively developed dissolution seams. Within the mineralized sections at XY, located within the Duo Lake Formation, S_1 forms spaced sulfide-rich dissolution seams that were previously interpreted as dewatering structures. These seams are up to 0.5 cm wide and show coarser-grained sphalerite, galena, and pyrite than the laminae they cut across. S_1 is visible at both the mesoscopic and microscopic scale, and its position relative to compositional layering (S_0) is a function of its location within a fold; however, at the outcrop scale S_0 has an enveloping surface that is always oblique to S_1 . It shows variable degrees of refraction across S_0 contacts, which is especially prevalent in the limbs of mesoscopic folds. A later kink fabric, S_k , is seen locally at a high angle to S_1 and may be genetically linked to movement on WNW- and NNE-striking faults that overprint the F_1 folds. Mapping constraints show that the faults exhibit normal dip-slip movement. Structural mapping shows that the geometry of the deposits is primarily controlled by folding and that contacts between units are stratigraphic in origin. S_1 and S_k are the only foliations mapped. No shear zone fabrics were identified and there is no evidence that bedding has been transposed.

Fig. 3. Geologic map of the 2020 XY region field area. Stereonet of all bedding data collected in mapping area is shown in the lower right (lower hemisphere projection, equal area). From Kamal and Hickey (2020).

