

# SEG 100 Conference: Celebrating a Century of Discovery

---

**ST.083**

## **Geology, Hydrothermal Alteration and Ore Geochemistry of Prospect-47, Alta Floresta Gold Province, Brazil: The First Insights into a New Cu ± Au ± Mo Prospect**

Danilo Pineschi, Rafael R. Assis

University of São Paulo, Institute of Geosciences, São Paulo, SP, Brazil

The Alta Floresta Gold Province, located in the southwestern sector of the Amazonian Craton (State of Mato Grosso, Brazil), is mainly composed of Paleoproterozoic plutonic, volcanic, and volcano-sedimentary units interpreted to have originated in magmatic arc to postorogenic settings from 2.1 to 1.7 Ga. Particularly in its eastern sector, relatively oxidized (magnetite-bearing) I-type, calc-alkaline to subalkaline, medium- to high-K, meta to peraluminous granitic rocks and volcanic sequences host a number of gold and polymetallic deposits. Gold usually occurs as inclusions in pyrite, the main ore mineral, with subordinate chalcopyrite, galena, sphalerite, molybdenite, hematite, and magnetite. Most gold-rich deposits are interpreted as similar to porphyry-epithermal systems. Despite being a primarily Au-rich province, several mining companies have put efforts into greenfield exploration campaigns aimed at discovering Cu occurrences. This is the case of Prospect-47, a new Cu ± Au ± Mo prospect in the eastern sector of the Province.

Logging of seven drill cores from Prospect-47 revealed that mineralization is hosted by (i) very fine- to fine-grained porphyritic quartz monzodiorite, (ii) medium- to coarse-grained quartz monzodiorite, (iii) fine- to coarse-grained porphyritic granodiorite (intrusive in the previous rock types), and (iv) quartz-feldspar porphyry of rhyolitic composition (intrusive in all rock types). These rock types were temporally subjected to (1) pervasive potassic alteration, characterized by K-feldspar, biotite, or both replacing igneous minerals; (2) pervasive sericitic alteration that overprints the potassic alteration and affects mostly igneous plagioclase; and (3) pervasive chloritic alteration characterized by hydrothermal chlorite, epidote, and carbonate overprinting previous hydrothermal alterations. All rock and hydrothermal alteration types are crosscut by syn- to late-mineralization quartz-carbonate veins and epidote- or chlorite-rich veinlets. Mineralization is mainly associated with potassic alteration zones and is characterized by disseminated pyrite, chalcopyrite, bornite, and molybdenite, but it also occurs within quartz-carbonate veins and epidote- and chlorite-rich veinlets. Hydrothermally brecciated zones are uncommon but also mineralized. Preliminary ore geochemistry data reveal Cu values ranging from <50 to 9500 ppm, Mo values ranging from <1 to 996 ppm, and Au values ranging from <0.01 to 0.17 ppm. The highest Cu, Au, and Mo values are usually related to potassic alteration zones in the granodiorite and the porphyry. Cu and Au values display a positive linear correlation, whereas Cu and Mo and Au and Mo show no correlation. Ag (<1 to 14 ppm), Zn (<1 to 1619 ppm), and Pb (<3 to 237 ppm) values show no correlation with Cu, Au, and Mo, nor with each other.

These preliminary results suggest that the prospect may be genetically related to a magmatic-hydrothermal system. Further work, such as (i) petrography of host rocks and ore zones; (ii) whole-rock geochemistry of host rocks; and (iii) zircon U-Pb (SHRIMP) geochronology will provide better insights into the geological, tectonic, and metallogenetic history of the prospect.

Fig. 1. Location map of Prospect-47 and nearby mineral deposits, Amazonian Craton.

