

# SEG 100 Conference: Celebrating a Century of Discovery

---

**ST.213**

## **Abrupt Switch in Magmatic Plumbing to Tap Porphyry Copper-Fertile Magmas**

Lawrence C. Carter<sup>1</sup>, Simon R. Tapster<sup>2</sup>, Ben J. Williamson<sup>1, 3</sup>, Yannick Buret<sup>3</sup>, David Selby<sup>4, 5</sup>, Daniel Parvaz<sup>6, 7</sup>

1. Camborne School Of Mines, University of Exeter, Cornwall, United Kingdom, 2. British Geological Survey, Nottingham, United Kingdom, 3. Natural History Museum, London, United Kingdom, 4. Durham University, Durham, United Kingdom, 5. State Key Laboratory of Geological Processes and Mineral Resources, Wuhan, China, 6. Selfrag AG, Kerzers, Switzerland, 7. Lightning Machines, Manchester, United Kingdom

Porphyry-type Cu deposits are amongst the most intensively studied ore systems (Richards, 2018) and will play a major role in the shift towards a green economy. They form from large, long-lived, multicomponent magmatic complexes, mostly in subduction-related settings (e.g., Seedorff et al., 2005; Sillitoe, 2010; Wilkinson, 2013). Due to limited or patchy exposure of their 3D architecture (Seedorff et al., 2008), it is often difficult to determine when and why a relatively small number of systems, and more specifically certain batches of magma within them, produce deposits. We address this from field, mineralogical, geochemical (whole-rock and zircon), and geochronological (zircon U-Pb CA-ID-TIMS and molybdenite Re-Os) studies of the unique, comparatively complete ~8-km-deep cross section through the archetypal Yerington porphyry system, Nevada (Dilles, 1987).

We show that long-lived (>1.5 m.y.), relatively shallow (~1-8 km; Dilles, 1987) emplacement of the premineralisation McLeod Hill quartz monzodiorite and Bear quartz monzonite plutons, from a mid-crustal source (~15-20 km depth), rapidly switched (over ~100 k.y.) to slightly deeper intrusion (~4-8 km; Dilles, 1987) of more evolved magmas from a lower crustal source (~20-40 km depth) to form the Luhr Hill granite. Inward crystallisation of the latter produced Cu- and volatile-rich magmas, which were emplaced upwards as multiple generations of aplite dikes. These acted as crystal mush conduits (Carter et al., 2021) for exsolving magmatic-hydrothermal fluids from large volumes of underlying magma, over a period of up to ~1.1 m.y., to form the four known porphyry Cu deposits within the apices of the Luhr Hill granite. This framework, where porphyry deposit formation relied on a rapid switch in magma source and evolution to more copper-fertile compositions and the focussing of fluids through crystal mush dikes (Carter et al., 2021), offers new constraints for exploration models.

### **References**

- Carter, L.C. et al., 2021, *Commun. Earth Environ.*, 2, 59.  
Dilles, J.H., 1987, *Econ. Geol.*, 82, 1750-1789.  
Richards, J.P., 2018, *Econ. Geol.*, 113, 1225-1233.  
Seedorff, E. et al., 2005, *Econ. Geol.*, 100, 251-298.  
Seedorff, E. et al., 2008, *Econ. Geol.*, 103, 939-956.  
Sillitoe, R.H., 2010, *Econ. Geol.*, 105, 3-41.  
Wilkinson, J.J., 2013, *Econ. Geol.*, 6, 917-925.