

SEG 2022 Conference: Minerals For Our Future

Lead Isotope and Fluid Inclusion Investigations of the Hicks Dome Critical Mineral Resource, Illinois-Kentucky Fluorspar district, USA

Mitchell M. Bennett¹, Albert H. Hofstra¹, Mike Pribil¹, Allen K. Andersen², Wayne Premo³, Anne E. McCafferty¹, Anton Maria⁴, Laurence M. Nuelle⁵

1. US Geological Survey, Denver, CO, USA, 2. US Geological Survey, Spokane, WA, USA, 3. US Geological Survey (Emeritus), Denver, CO, USA, 4. University of Southern Indiana, Evansville, IN, USA, 5. Hicks Dome LLC, Hardin County, IL, USA

Deep drilling at Hicks Dome has defined an inferred critical mineral resource of 65.8 Mt with average grades of 15.8% fluorite, 4.8% barite, 1.3% TiO₂, 0.3% REY oxide, 0.2% Nb₂O₅, and 0.16% BeO using a fluorite cut-off grade of 10%. Fine-grained purple fluorite mineralization occurs in solution collapse breccias with a rock flour-matrix. Breccias are hosted in Ordovician to Silurian platform carbonate rocks at depths of 700-1000 m. The resource is associated with 271 Ma lamprophyre dikes and diatremes, an upper crustal magnetic anomaly, and a Th radiometric anomaly at the surface. It is hypothesized the origin for the dome and the HREE enrichment is the product of fluid exsolved from a concealed carbonatite intrusion.

SEM-EDS raw x-ray intensity maps and laser Raman spectroscopy were utilized to identify mineral phases in high grade ore. X-ray intensity maps were grouped with Aztec algorithms to form phase maps that identify fluorite, barite, Pb-Zn-Fe-sulfides, Ca-Mg±Sr±Ba carbonates, apatite, Th-REE phosphates, and Nb-Ti oxides. In-situ Pb isotope analyses of ore and gangue minerals by LA-MC-ICP-MS is proceeding to establish an isochron and to constrain the source(s) of Pb in the system.

Quartz and fluorite contain distinct secondary fluid inclusion populations. Two-phase liquid-rich inclusions are irregular to equant in shape, with a small to medium bubble. Three-phase liquid-vapor-solid inclusions are morphologically similar to the liquid-rich inclusions. Sparse vapor-rich inclusions occur with liquid-vapor inclusions in fluorite. Three-phase water-CO₂ inclusions are restricted to quartz, contain varying ratios of vapor- and liquid-CO₂ and liquid water; some assemblages contain a solid daughter mineral. Pyrite mineral inclusions are restricted to late-stage fluorite, while primary liquid-rich inclusions are present in calcite and late-stage fluorite. Microthermometric and laser Raman characterization of these fluid inclusion populations is underway to constrain the temperature, salinity, and gas compositions of hydrothermal fluids in the system.