

## **Alteration Geothermometry and Exploration Implications of the Gajah Tidur Porphyry System, One of Two Systems that Formed the Supergiant Grasberg Cu-Au-(Mo) Deposit**

Reza Al Furqan<sup>1</sup>, Antonio Arribas<sup>2</sup>, Clyde Leys<sup>3</sup>, Tetsumaru Itaya<sup>4</sup>, Adi Sulaksono<sup>5</sup>, Yasushi Watanabe<sup>1</sup>  
1. Akita University, Akita, Japan, 2. University of Texas at El Paso, El Paso, TX, USA, 3. Eksplorasi Nusa Jaya (Affiliate of Freeport-McMoran Inc), Jakarta, Indonesia, 4. Japan Geochronology Network, Okayama, Japan, 5. UPN Veteran Yogyakarta, Yogyakarta, Indonesia

Based on geochronological data and field relationships, researchers have recently proposed that the Grasberg Cu-Au-(Mo) deposit comprises two overlapping porphyry Cu systems, the deeper and older (~3.4 Ma) Gajah Tidur (GT) Cu-Mo-(Au) system and the shallower and younger (~3.1 Ma) Main Grasberg (MG) Cu-Au system. Dating of abundant hydrothermal muscovite and subordinate biotite in this study supports the age relationships between the two porphyry systems and indicates that alteration and associated Cu-Mo-(Au) mineralization beneath the now mined-out Grasberg open pit is mainly associated with the GT porphyry system. As well as a ~1-km difference in emplacement depth compared to MG, a different metal signature (Cu-Mo-(Au) vs. Cu-Au), and distinct parent intrusion geometry, Cu mineralization in GT is mainly associated with a lower-temperature alteration assemblage. Sulfur isotope temperatures from sulfate-sulfide pairs show that initial GT magmatic-hydrothermal fluids formed K-feldspar-biotite alteration at temperatures up to 590°C. Cooling of this fluid from ~460° to ~370°C resulted in white mica-dominated alteration associated with anhydrite, chlorite, chalcopyrite, and pyrite. Further cooling (340°–280°C), outward from the core of the hydrothermal system, is characterized by a muscovite-quartz ± pyrophyllite assemblage associated with covellite-pyrite mineralization. While Cu mineralization in the GT porphyry system is largely associated with white mica assemblages (muscovite-anhydrite-chlorite and muscovite-quartz ± pyrophyllite), high-grade Cu and Au mineralization in the MG system is chiefly associated with a K-feldspar-biotite-magnetite assemblage formed at higher temperature than the GT K-feldspar-biotite alteration. The overlapping GT and MG porphyry systems formed the supergiant Grasberg Cu-Au-(Mo) porphyry deposit, within which GT contributed an estimated 30% for Cu, 10% for Au, and 75% for Mo. The existence of two closely spaced, well-mineralized porphyry systems where the older system was emplaced at a greater depth than the younger one is an unusual occurrence that has significant exploration implications.