

## Contrasting Tectonomagmatic Conditions for Porphyry Cu and Iron Oxide-Apatite Deposits in Phanerozoic Arcs

Xuyang Meng<sup>1</sup>, Jingwen Mao<sup>1</sup>, Adam Simon<sup>2</sup>, Chao Duan<sup>3</sup>, Guiqing Xie<sup>1</sup>, Huimin Su<sup>4</sup>, Tong Hou<sup>1</sup>, Ke Shi<sup>5</sup>, Nian Chen<sup>1</sup>

1. China University of Geosciences (Beijing), Beijing, China, 2. University of Michigan, Ann Arbor, MI, USA, 3. Institute of Mineral Resources, Chinese Academy of Geological Sciences (CAGS), Beijing, China, 4. China University of Geosciences (Wuhan), Wuhan, China, 5. MNR Innovative Center of Coverage Area Deep Resource Exploration Engineering Technology, Geological Survey of Anhui Province, Hefei, China

Porphyry Cu  $\pm$  Mo  $\pm$  Au and iron oxide ( $\pm$  apatite, IOA;  $\pm$  copper-gold, IOCG) deposits commonly occur in close spatial and temporal proximity in Phanerozoic arc-related settings. However, the evolution of these deposit types in an evolving arc setting remains poorly understood. Here, we systematically evaluated the P-T- $fO_2$  conditions and S-Cl contents for the dioritic to granodioritic source magmas for representative magmatic-hydrothermal porphyry-skarn Cu  $\pm$  Au (150–135 Ma) and IOA (~130 Ma) deposits that formed under transpressional and transtensional settings in the Middle-Lower Yangtze River Metallogenic Belt, China. Our estimates show that, compared to IOA deposits, the porphyry-skarn-related magmas were relatively felsic, cooler, more hydrous, and emplaced relatively deeper ( $264 \pm 23$  vs.  $143 \pm 17$  MPa for IOA-related magmas; 2SE). Apatite data reveal that the porphyry-related and IOA-related magmas had comparable pre-degassed S concentrations ( $\sim 0.13 \pm 0.06$  vs.  $\sim 0.16 \pm 0.09$  wt. % on average), but that IOA-related magmas had higher pre-degassed Cl/H<sub>2</sub>O ratios ( $\sim 0.10$ – $0.15$  vs.  $0.02$ – $0.08$  wt. % for porphyry-related magmas) that decreased by a factor of two after magma degassing. Magmatic  $fO_2$  estimated using zircon and amphibole, reported in log units relative to fayalite-magnetite-quartz (FMQ) redox buffer, gradually increased during cooling of the porphyry-skarn-related magmas ( $\Delta FMQ +0.7$  to  $+2.5$ ) at  $950^\circ$  to  $800^\circ C$  and decreased to  $\Delta FMQ +1$  at  $700^\circ C$ . In contrast, the magmatic  $fO_2$  for the IOA-related magmas varied significantly from  $\Delta FMQ -1.5$  to  $\Delta FMQ +2.5$  but generally showed an increasing trend with cooling from  $970^\circ$  to  $700^\circ C$ . We propose that the porphyry-skarn Cu deposits in the MLYRMB formed in a compressional setting in response to normal plate subduction that favors storage and evolution of S-rich hydrous ore-forming magmas at variable crustal levels, whereas the back-arc extensional setting that formed because of slab rollback led to rapid degassing of the Cl-rich magmas.