

Melt Inclusion Constraints on the Initial Metal Budget of Suspected Causative Magmas in the Au-Rich Doyon-Bousquet-LaRonde Mining District, Abitibi Greenstone Belt, Québec

Kevin Neyedley¹, Jacob J. Hanley¹, Patrick Mercier-Langevin², Zoltan Zajacz³, Alexandra Tsay³

1. Department of Geology, Saint Mary's University, Halifax, NS, Canada, 2. Geological Survey of Canada, Quebec, QC, Canada, 3. Department of Earth Sciences, University of Geneva, Geneva, Switzerland

The synvolcanic Mooshla Intrusive Complex (MIC) in the Doyon-Bousquet-LaRonde mining camp, Abitibi greenstone belt, Canada, may be genetically associated with numerous Au-rich VMS, epizonal intrusion-related Au ± Cu, and shear zone-hosted orogenic Au and/or remobilized VMS deposits. To understand the magmatic evolution of the MIC and its relationship to these Au-rich deposits, analysis of silicate melt inclusions (SMIs) preserved in zircon and quartz provides a means to determine the original major- and trace-element composition and metal tenor of the magmatic system. The use of SMIs is preferred for quantification of the pre-emplacement melt chemistry (in particular, ore metals) compared to bulk-rock compositions commonly modified by magmatic (e.g., degassing) and post-solidus processes.

Characterization of SMI chemistry from different lithologies of the MIC (gabbro, quartz diorites, tonalites, and trondhjemites) was done by integrating petrography with SEM-EDS, EMPA, and LA-ICP-MS analyses. Data presented constitute the first SMI study in Archean-age rocks within Canada.

Melt inclusions, trapped at depths of ~18.5 to 40 km (based on thermobarometry in a parallel study) are sub-alkaline to alkaline and rhyolitic in composition from all lithologies. Importantly, within individual lithologies, SMIs span from sub-alkaline to alkaline compositions, indicating melt entrapment at various stages of magmatic evolution, requiring contamination and/or magma mixing. Present in both zircon- and quartz-hosted SMIs are accidentally trapped sulphide phases (enriched in Au, Ag, Cu, Bi), indicating the melts were sulphide-saturated at the time of SMI entrapment. In quartz-hosted SMIs with no accidentally trapped sulphide, Au concentrations range from 10 to 120 ppb (70 ± 50 ppb; $n = 6$). Enrichment of SMIs in Au and other metals (e.g., Ag, Cu, Sb, Bi) typical of porphyry-epithermal environments in arc settings reinforces that consideration must be made for the direct involvement of magmatically sourced metals transferred into shallow ore-forming hydrothermal systems through deep magmatic devolatilization.