

Factors for Classification of the Precious and Base-Metal Mineralization in Epithermal Deposits: Evidence from Apatite Chemistry in Ore-Related Igneous Activity

Masaya Koseki¹, Shota Satori², Yasushi Watanabe¹, Takuya Echigo¹, Shogo Aoki¹

1. Akita University, Graduate School of International Resource Sciences, Akita, Japan, 2. National Institute of Advanced Industrial Science and Technology (AIST), Geological Survey of Japan, Tsukuba, Japan

Metal assemblages concentrated in epithermal deposits are related to Cl and S contents in hydrothermal fluids. The differences of Cl and S contents in hydrothermal fluids may reflect the difference of ore-related magma compositions. We estimated Cl and S contents exsolved from magmas by using apatite chemistry in the felsic rocks associated with the Miocene Kitano-o Au-Ag and Inaushi Cu-Pb-Zn deposits in northeastern Hokkaido, Japan.

The EPMA analysis indicated that Cl content in the apatite is 0.56–0.97 wt% in the Inaushi dacite, while it is 0.37–0.45 wt% in the Kitano-o rhyolite. Based on the partition coefficients between apatite and melt, the calculated maximum Cl contents in the Inaushi and Kitano-o melts are 1.48 wt% and 0.96 wt%, respectively. Both apatites contain ~100 ppm S, indicating <10 ppm of S in the Inaushi and Kitano-o melts.

We also traced the Cl in the melt during magmatic fractionation of the Inaushi dacite by using apatites of different crystallizing stages, which are determined by paragenetic sequence of apatite-hosting minerals. Maximum Cl content is 0.78 wt% during the early plagioclase crystallization and 0.97 wt% during the amphibole crystallization, and then it decreases to 0.87 wt% and 0.83 wt% during titanomagnetite and groundmass crystallization, respectively.

The significant difference of Cl content and similar small S content between the Inaushi dacitic and Kitano-o rhyolitic melts estimated by apatite chemistry suggest that the Cl content in the epithermal fluids for the Inaushi base-metal and Kitano-o precious metal mineralization was inherited from the those in the magmas associated with these deposits. At Inaushi, Cl content was estimated to have exsolved from melt during amphibole-titanomagnetite crystallization stages.